




Principle of Quantum Cascade Laser




New type of structure



Laser characteristics at B = 0T



Use of magnetic field - experimental results



Interpretation


Interaction electron-interface roughness (Elastic)

Calculated upper state lifetime considering the relaxation via the interface roughness scattering from $|9,0\rangle$ Landau level to the Landau levels of all the subbands of the extraction miniband.


$$E_{9,0} + \frac{1}{2} \hbar \omega_c = E_{7,0} + \left(m + \frac{1}{2}\right) \hbar \omega_c$$

with $m = 1, 2, 3, \dots$

$$\frac{1}{\tau_{9,0}^{\text{roughness}}} = \sum_n \left(\frac{1}{\tau_{9,0-7,n}^{\text{roughness}}} + \sum_{i=3}^6 \frac{1}{\tau_{9,0-i,n}^{\text{roughness}}} \right)$$



Interaction electron-LO phonon (Inelastic)



$$E_{9,0} + \left(n + \frac{1}{2}\right) \hbar \omega_c = E_{7,0} + \frac{1}{2} \hbar \omega_c + \hbar \omega_{\text{LO}}$$

with $n = 1, 2, 3, \dots$

This series of oscillations implies a hot electrons population injected in Landau levels $|9, n\rangle$. ($T_e = 150\text{K}$)

Using a complete set of rate equations for levels $|9\rangle$ and $|7\rangle$, we demonstrate this non radiative mechanism breaks inversion population.

Conclusion

- Two non-radiative mechanisms are pointed out in this sample
- Demonstration of high electronic temperature
- Need to lower electronic temperature in QCL by new design